## CLAIMS

1. A compound of the formula:-

or a pharmaceutically acceptable salt thereof, wherein R is phenyl substituted by 1 to 3 substituents each independently selected from halo CF3 and OCF4.

 $R^1$  is  $C_1 - C_4$  alkyl;


R<sup>2</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl:

A X is GH or N; and

Y is F or Cl.

- 2. A compound of claim 1, wherein R is phenyl substituted by 1 or 2 substituents each independently selected from fluoro and chloro.
- 3. A compound of claim 2, wherein R is 2-fluorophenyl, 2,4-diffluorophenyl, 2-chlorophenyl, or 2,4-dichlorophenyl.
  - 4. A compound of claim 3, wherein R<sup>1</sup> is methyl.
- 5. A compound of claim 4, wherein R<sup>2</sup> is hydrogen or methyl.
- 6. The compound of claim 5, wherein R<sup>2</sup> is hydrogen, R is 2,4 diffuorophenyl, X is N and Y is fluoro.
  - 7. The compound of claim 5, wherein R<sup>2</sup> is hydrogen, R is 2,4-difluorophenyl, X is CH and Y is fluoro.

8. The compound of claim 5, of the formula



where R<sup>2</sup> is hydrogen, R is 2,4-difluorophenyl, X is N and Y is fluoro.

9. The compound of claim 5, of the formula

wherein R<sup>2</sup> is hydrogen, R is 2,4-difluorophenyl, X is CH and Y is fluoro.

10. A method of treating a fungal infection in a mammal which comprises administering to said mammal an antifungal effective amount of a compound of the formula

or a pharmaceutically acceptable salt thereof, wherein R is phenyl substituted by 1, to 3 substituents each independently selected from halo er 3 and ocr 3;

 $R^{1}$  is  $C_{1}-C_{4}$  alkyl;  $\chi$  is CH or N; and Y is F or  $C_{1}$ .

- 11. A method of claim 10, wherein R is phenyl substituted by 1 or 2 substituents each independently selected from fluoro and chloro.
- 12. A method of claim 11, wherein R is 2-fluorophenyl, 2,4-difluorophenyl, 2-chlorophenyl or 2,4-dichlorophenyl.
  - 13. A method of claim 12, wherein R1 is methyl.
- 14. A method of claim 13, wherein R<sup>2</sup> is hydrogen or methyl.
- 15. A method of claim 14, wherein R<sup>2</sup> is hydrogen, R is 2,4-difluoropheny, X is N or CH and Y is fluoro.
  - 16. A method of claim 14, wherein the compound has the formula

CH<sub>3</sub>
HO
R
Y
N
R
Y
N
R

wherein R is 2,4-difluorophenyl, X is N or CH and Y is fluoro.

- 17. A method of claim 16, wherein the fungal infection is caused by the Aspergillus sp fungi.
- 18. A method of claim 17, wherein the Aspergillus sp fungi is Aspergillus fumigatus.
- 19. A pharmaceutical composition in unit dosage
  form for treating a fungal infection in a human which
  comprises a compound according to claim 1, together
  with a pharmaceutically acceptable diluent or carrier.
  - 20. A pharmaceutical composition of claim 19, wherein the compound according to claim 1 is complexed with a hydroxyalkyl derivative of a cyclodextrin.

21. A pharmaceutical composition of claim 20, wherein the hydroxyalkyl derivative is a hydroxypropyl derivative and the cyclodextrin is alpha- or beta-cyclodextrin.

22. A compound of the formula

$$\begin{array}{c|c}
R^1 & R^2 \\
\downarrow & & \\
R & Y
\end{array}$$
or
$$\begin{array}{c|c}
R^1 & R^2 \\
\downarrow & \\
R & Y
\end{array}$$

wherein R is phenyl substituted by 1 to 3 substituents each independently selected from halo, -CF<sub>3</sub> or -OCF<sub>3</sub>;

R<sup>1</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl; R<sup>2</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl; X is CH or N;

Y is Cl or F; and

z is chloro, bromo or  $C_1-C_4$  alkylsulfonyloxy.

23. A compound of the formula

wherein R is phenyl substituted by 1 to 3 substituents each independently selected from halo, -CF<sub>3</sub> and -OCF<sub>3</sub>;

 $z^2$  and  $z^3$  are each independently selected from H and a group that may be selectively removed by reduction, with the proviso that  $z^2$  and  $z^3$  cannot both be H.

24. A compound in claim 23 wherein the group that may be selectively removed by reduction is halo.

25. A compound in claim 24 wherein  $z^2$  is chloro and  $z^3$  is H.

add /